

## Haploinsufficiency of DNA damage response genes and their potential influence in human genomic disorders

Article (Accepted Version)

O'Driscoll, Mark (2008) Haploinsufficiency of DNA damage response genes and their potential influence in human genomic disorders. *Current Genomics*, 9 (3). pp. 137-146. ISSN 1389-2029

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/29179/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Haploinsufficiency of DNA Damage response genes and their  
potential influence in human genomic disorders.**

**Mark O'Driscoll**

**Genome Damage & Stability Centre**

**University of Sussex**

**Falmer**

**Brighton**

**East Sussex**

**BN1 9RQ**

**UK**

**Ph. 0044 (0) 1273 877 515**

**Fax. 0044 (0) 1273 678 121**

**[m.o-driscoll@sussex.ac.uk](mailto:m.o-driscoll@sussex.ac.uk)**

**Keywords : DNA damage response, ATR, haploinsufficiency, genomic disorders.**

## **Abstract.**

Genomic disorders are a clinically diverse group of conditions caused by gain, loss or re-orientation of a genomic region containing dosage-sensitive genes. One class of genomic disorder is caused by hemizygous deletions resulting in haploinsufficiency of a single or, more usually, several genes. For example, the heterozygous contiguous gene deletion on chromosome 22q11.2 causing DiGeorge syndrome involves at least 20-30 genes. Determining how copy number variation (CNV) affects human variation and contributes to the aetiology and progression of various genomic disorders represent important questions for the future. Here, I will discuss the functional significance of one form of CNV, haploinsufficiency (ie. loss of a gene copy), of DNA damage response components and its association with certain genomic disorders. There is increasing evidence that haploinsufficiency for certain genes encoding key players in the cells response to DNA damage has a functional impact, particularly those of the Ataxia Telangiectasia and Rad3-related (ATR)-pathway. I will review this evidence and present examples of some well known clinically similar genomic disorders that have recently been shown to be defective in the ATR-dependent DNA damage response. Finally, I will discuss the potential implications of a haploinsufficiency-induced defective DNA damage response for the clinical management of certain human genomic disorders.

## **1. Introduction.**

It has long been appreciated that changes in gene copy number are associated with phenotypes in humans. Perhaps the most well know example of this is the

trisomy 21 causative of Down syndrome. Here, increased expression of the genes on chromosome 21 results directly or indirectly in a clinically heterogeneous disorder incorporating cognitive impairment, facial dysmorphology, growth retardation, cancer predisposition, microcephaly, heart and skeletal abnormalities [1]. Interestingly, it was recently shown that CNV is in fact a common genetic trait in clinically unaffected or 'normal' individuals [2]. Indeed, the first complete genomic sequence from an individual, the so-called 'Venter genome' yielded a surprising level of CNV, highlighting the plasticity of the human genome [3]. The phenomenal recent revolution in the sensitivity and widespread usage of array-based Comparative Genomic Hybridization (array-CGH) techniques has led to CNV being described as the 'Breakthrough of the Year' by the journal Science [4]. The widespread use of a-CGH has facilitated the description of several novel genomic disorders and aided in the detailed genetic characterisation of known genomic disorders. A persistent challenge to clinical geneticists and researchers is to unravel exactly how changes in CNV of specific genes or various combinations of genes can impact on normal development. These issues have recently been extensively reviewed elsewhere and the reader is directed to these sources for an overview of a-CGH technology and its applications [5-12]. Here, I will discuss the potential role of haploinsufficiency of DNA damage response components in genomic disorders. Contiguous gene deletion disorders represent a clinically diverse group of human genomic disorders caused by distinct heterozygous chromosomal deletions usually involving several genes [5,13]. It is assumed that the various clinical manifestations of different genomic disorders arise from the combined impact of haploinsufficiency of multiple genes [14]. It is likely that there are critical genes or pathways sensitive to haploinsufficiency either alone or when combined with haploinsufficiency of other genes. There is increasing

evidence from both human and murine studies suggestive of a cellular impact of haploinsufficiency of genes that control different aspects of the response to DNA damage. Since our genomes are constantly exposed to exogenously-derived (eg. UV radiation) and endogenously-derived (eg. metabolically generated reactive oxygen species) DNA damaging agents, an impaired ability to detect and/or respond appropriately to these threats can impact on the maintenance of genetic stability. There are many examples of human Mendelian disorders defective in the repair of or response to DNA damage [15]. The importance of these pathways is demonstrated by the increase in cancer predisposition and developmental abnormalities associated with these conditions [15]. An important DNA damage response (DDR) pathway that appears to be affected by haploinsufficiency is the ATR-dependent DDR (ATR-DDR).

## **2. The ATR-dependent DNA damage response (ATR-DDR).**

The DDR can be divided into DNA repair processes and signal transduction processes that sense DNA damage and co-ordinate the appropriate response such as cell cycle checkpoint activation, DNA repair and/or apoptosis. Two phosphoinositol-3-kinase-like protein kinases (PIKK) co-ordinate the signal transduction response to DNA damage in mammalian cells [16-18]. Ataxia Telangiectasia Mutated (ATM) is activated following DNA double strand breaks (DSBs) whilst a related kinase, Ataxia Telangiectasia and Rad3-related (ATR), responds to single stranded regions of DNA. Single stranded DNA (ssDNA) can occur as an intermediate during the activity of several excision-mediated repair pathways (nucleotide excision repair and base excision repair), during uncoupling of the transcriptional machinery from its DNA

template or similarly following stalling of the DNA replication machinery at DNA lesions or single strand breaks. Following the exposure of ssDNA it is quickly coated by RPA a heterotrimeric protein complex (RPA1-3) that plays a role in many aspects of DNA metabolism (eg. replication, repair, transcription). ATR is recruited to the ssDNA via its binding partner ATRIP (ATR Interacting Protein) [19,20].

Topoisomerase binding protein 1 (TopBP1) appears to be required for optimal ATR kinase activity [21]. Phosphorylation of the histone H2A variant H2AX on Ser-139 (called  $\gamma$ -H2AX) is one of the earliest detectable PIKK-dependent responses to DNA damage and is required for the retention of additional damage response proteins at the damage site [22-24]. The Mre11/Rad50/NBS1 complex, the Rad17/Rfc2-5 and Rad9/Rad1/Hus1 complexes are recruited to the site of damage independently of ATR/ATRIP and are also phosphorylated by ATR [25,26]. Retention of these complexes facilitates ATR's ability to phosphorylate downstream substrates including Brca1, p53 and its effector kinase Chk1 [15].

There is a large amount of functional overlap between ATM and ATR. In fact, both kinase's phosphorylate mainly the same substrates in response to DNA damage (eg. Mre11/Rad50/NBS1 complex, p53, Brca1). DSBs can undergo exonucleolytic resection generating ssDNA overhangs, hence generating an ATR activating substrate [27]. Conversely, ssDNA generated at stalled replication forks can collapse producing overt DSBs, hence generating an ATM activating substrate [28]. Nevertheless, congenital defects in ATM and ATR are associated with clinically distinct human disorders. Mutations in *ATM* cause Ataxia telangiectasia, a progressive degenerative neurological disorder characterised by ataxia and lymphoid malignancy predisposition

[29,30]. Mutations in *ATR* result in Seckel syndrome a disorder characterised by microcephaly and growth retardation [31,32].

### **3. ATR and genomic stability.**

ATR is required to maintain genomic stability. Whilst it has a role in the stabilisation of stalled replication forks, the absence of ATR results in a very specific type of genomic instability, namely DNA Fragile Site expression. DNA Fragile Sites (DFS) are large (>100Kb) distinct genomic regions that exhibit breaks under conditions of replicative ‘stress’ [33,34]. Such stresses can be induced in the laboratory using DNA replication inhibitors such as aphidicolin or by folate deficiency, which can indirectly impact on the availability of dNTP’s. There are about 75-80 DFS throughout the human genome but it is not really clear why DFS are so unstable. Most tend to be relatively AT-rich and contain more areas of flexibility than non-fragile site regions. Studies on the replication timing of certain fragile sites (FRA3B, FRA7H, FRAXA on chromosome 3, 7 and X respectively) indicate that they are replicated very slowly [33]. DFS are ‘hot spots’ for sister chromatid exchanges (SCE’s) and are also thought to play a role in gene amplification events via a breakage-fusion-bridge cycle [33]. Breakage at or ‘expression of’ DFS is associated with many cancers. For example, the *FHIT* tumour suppressor gene spans the DFS FRA3B [35]. This gene is often re-arranged or partially deleted in a wide range of tumours (lung, ovarian, breast, esophageal) [36]. Pioneering work by the Glover laboratory identified ATR as the first protein required to mediate DFS stability [37]. Subsequently, several other proteins have been implicated in maintaining DFS stability including the Brca1, SMC1, WRN helicase, Chk1 and the Fanconi anaemia

pathway components FANC-A, FANC-B and FANC-D2 [33,38-41]. Interestingly, most of these proteins are known direct ATR substrates.

#### **4. Seckel syndrome: an ATR-pathway defective disorder.**

Seckel syndrome, originally described as 'Bird headed Dwarfism' in a 1960 monograph by Dr. Helmut Seckel, is a disorder characterised by severe microcephaly, isolated skeletal abnormalities (clinodactyly, thoracic kyphosis, ivory epiphesis) and a dramatic proportionate primordial dwarfism [42]. Microcephaly is a clinical term describing a reduction in occipitofrontal (or head) circumference greater than 3 standard deviations (-3 s.d.) below the age-related mean. This reduced head circumference is a consequence of premature closure and fusion of the cranial sutures reflecting the underlying reduction in brain volume. The aetiology of microcephaly is complex. It can occur in the context of genetic (e.g. syndromal) or non-genetic (e.g. intrauterine infection) situations. Microcephaly is particularly pronounced in Seckel syndrome [31,42,43]. The first genetic defect associated with Seckel syndrome was described by in 2003. A single synonymous hypomorphic mutation in *ATR* was identified in five affected individuals in two consanguineous Pakistani families [31]. The mutation (A21201G) was shown to adversely impact on splicing resulting in dramatically reduced ATR expression in cells derived from the affected patients. Gene targeting of *ATR* in the murine system results in early embryonic lethality [44,45]. Hence, these ATR-defective Seckel syndrome cells (ATR-S) proved a useful tractable model to investigate ATR-pathway function in the mammalian setting. ATR-S cells exhibited a diminished ability to phosphorylate ATR substrates following DNA damage-induced ATR-pathway activation (e.g.  $\gamma$ H2AX formation and p53-



serine-15 phosphorylation) as well as defective cell cycle checkpoint arrest and increased DFS expression [31,46,47].

Seckel syndrome is known to be genetically heterogeneous [43,46,48-50]. Interestingly, Alderton *et al* showed that several unrelated non-*ATR* mutated Seckel syndrome cell lines all exhibited defective ATR-pathway function (e.g. failure of ATR-dependent G2-M cell cycle checkpoint arrest) [46]. Hence, in agreement with the known genetic heterogeneity of this condition, whilst not all Seckel syndrome cases are caused by mutations in *ATR* itself, interestingly, all Seckel syndrome cell lines (*ATR*-S and non-*ATR*-S) exhibit compromised ATR-pathway function [46]. A further interesting cellular feature of ATR-pathway defective Seckel cells is supernumerary centrosomes in 10-20% of mitotic cells [46]. A normal mitotic cell must only have two centrosomes that nucleate the microtubule spindles facilitating transfer of an equal and identical chromosome complement to the daughter cells [51,52]. The molecular aetiology of supernumerary mitotic centrosomes in Seckel syndrome cells is currently unclear. Centrosome orientation is fundamental for determining symmetric and asymmetric stem cell division in the embryonic neuroepithelium, an essential process for normal brain development [53-55]. Hence, this cellular feature may be a relevant contributor to the severe microcephaly characteristic of Seckel syndrome. Recently, the second genetic defect associated with Seckel syndrome was described. Mutations in *pericentrin (PCNT2)/kentrin* which encodes a structural centrosomal protein, were identified in several Seckel syndrome patients all of which exhibited defective ATR-pathway function [56]. This exciting finding further illustrates a functional link between the DDR, the cell cycle machinery and the centrosome. Furthermore, mutations in *PCNT2* were also recently found in

several microcephalic osteodysplastic primordial dwarfism type II (MOPD II) patients [57,58]. Whether *PCNT2*-mutated Seckel syndrome and MOPD II represent distinct disorders or allelic variants of the same condition remains an open question.

## **5. Other disorders that exhibit compromised ATR-pathway function.**

Compromised ATR-pathway function does not appear to be uniquely associated with Seckel syndrome. Several other known DDR disorders have been shown to be defective in aspects of ATR-pathway activity, particularly cell cycle checkpoint activation. Interestingly, whilst these disorders are all characterised by a distinct set of clinical features there is significant clinical overlap with Seckel syndrome, particularly concerning the developmental abnormalities such as microcephaly and growth retardation.

### ***5.1 Nijmegen breakage syndrome (NBS).***

NBS is caused by hypomorphic mutations in the *Nbs1* component of the Mre11/Rad50/NBS (MRN) complex [59,60]. The MRN complex plays a central role in ATM signalling where it is thought to recruit ATM to the DSB and also facilitate its ability to phosphorylate substrates such as p53 and Chk2 [61]. In fact, all the components of this complex are ATM substrates. Historically, NBS has been described as an A-T-like disorder as cell lines from both conditions exhibit radio-sensitivity and similar cell cycle checkpoint defects in response to DSBs. But, NBS1 is also an ATR substrate and NBS patients exhibit microcephaly and growth retardation, clinical features associated with Seckel syndrome and not A-T. Recently,

NBS cells were shown to be compromised for ATR-dependent checkpoint activation and other aspects of ATR-pathway function [26]. Therefore, NBS represents a human condition defective in elements of both ATM and ATR-pathway activity.

### ***5.2 Fanconi anaemia (FA).***

FA, is a genetically heterogeneous condition characterised by a progressive aplastic anaemia, skeletal abnormalities, microcephaly and lymphoid malignancy. FA is caused by mutations in different genes whose products co-ordinately function in the cellular response to DNA cross-links [62,63]. Several of these genes encode products that together mediate the monoubiquitylation and activation of the FANCD2 protein in response to DNA damage, particularly during S phase [62]. ATR and NBS1 have been shown to be required for this specific modification [64-66]. In fact, ATR-S and NBS patient-derived cell lines fail to monoubiquitylate FANCD2 following treatment with replication fork inhibitors [26]. In addition to undergoing monoubiquitylation, FANCD2 has been shown to be phosphorylated by ATM and ATR further highlighting the over-lap between these DDR pathways [26,64,67].

### ***5.3 MCPH1-dependent Primary Microcephaly.***

Autosomal recessive Primary Microcephaly, clinically characterised by the presence of a severe microcephaly in the absence of other overt clinical features, is a genetically heterogeneous condition composed of six distinct genetic complementation groups. To date, mutations in four genes, all of which encode centrosomal proteins, have been described for this disorder (*MCPH1*, *ASPM*,

*CDK5/RAP2, CENPJ*) [55,68-71]. *Microcephalin (MCPH1)*, the first Primary Microcephaly gene identified encodes a BRCT-containing product that has been implicated in the response to DSBs [72,73]. Importantly, work using MCPH1-patient derived cell lines with hypomorphic mutations in *MCPH1* indicated that these cells are defective in ATR-dependent checkpoint activation and also exhibit supernumerary mitotic centrosomes [74]. Furthermore, MCPH1 was shown to interact with Chk1, a substrate and downstream effector of ATR [74].

## **6. The impact of *ATR* haploinsufficiency: Murine studies.**

All of the disorders described so far that collectively exhibit defective ATR-pathway function are all caused by gene mutations inherited in an autosomal recessive manner. As mentioned earlier, a knockout mouse model for ATR does not exist due to early embryonic lethality of  $ATR^{-/-}$  blastocysts [44,45]. Whilst  $ATR^{+/-}$  animals are viable, re-evaluation of this murine work suggests that the ATR-DDR pathway is sensitive to gene copy number variation. Firstly,  $ATR^{+/-}$  mice are not born at the expected Mendelian frequency suggesting a requirement for a full diploid complement of ATR protein during normal development [75]. Furthermore, decreased survival and increased tumour incidence was recorded in the  $ATR^{+/-}$  animals compared to their  $ATR^{+/+}$  counterparts [75]. This is distinct to that of  $ATM^{+/-}$  embryos and mice. Heterozygous mutations in *ATR* have been observed in microsatellite unstable human colorectal carcinomas further suggesting that *ATR* haploinsufficiency may play a role in tumourigenesis [76]. In fact, it has been suggested that ATR acts as a haploinsufficient tumour suppressor under certain circumstances. Fang and colleagues showed that  $ATR^{+/-}$  mice when crossed into a mismatch repair defective

(*Mlh1*<sup>-/-</sup>) background (generating *ATR*<sup>+/-</sup>/*Mlh1*<sup>-/-</sup>) were highly susceptible to embryonic lethality and premature tumour development [77].

## **7. The impact of *ATR* haploinsufficiency: Human studies, including the association of *ATR*<sup>+/-</sup> with a genomic disorder.**

Fang and colleagues also showed that haploinsufficiency of *ATR* in human cells results in a clear DDR-defective phenotype [77]. Following gene targeting of *ATR* in the human HCT116 colorectal carcinoma cell system, they found a significantly increased expression of DFS as well as other gross chromosomal rearrangements and amplifications. Furthermore, these *ATR*<sup>+/-</sup> cells exhibited diminished *ATR*-mediated phosphorylation of its effector kinase Chk1.

Work from de Ru and colleagues along with that of O'Driscoll and colleagues has provided evidence for a cellular impact of *ATR* haploinsufficiency associated with a human genomic disorder [78,79]. Blepharophimosis-ptosis-epicanthus inversus syndrome (BPES) is a disorder characterised by a reduction in the dimensions of the palpebral fissures or eye sockets (blepharophimosis), drooping eyelids (ptosis) and inverted skin folds originating from the lower eyelids (epicanthus inversus) [80]. A variant of this disorder is also associated with ovarian failure and female infertility [81]. This disorder is caused by autosomal dominant mutations in or heterozygous deletion of *FOXL2*, a putative forkhead transcription factor [80]. In a review of the literature, de Ru and colleagues noted that most of the BPES patients with a cytologically detectable deletion or a microdeletion on chromosome 3q, where *FOXL2* resides, also exhibited microcephaly and short stature [78]. These are clinical

features not typically associated with ‘classical’ BPES. Indeed, it had been previously proposed that a putative gene for microcephaly was located close to the BPES-causative gene on chromosome 3q (reviewed in [78]). De Rue and colleagues mapped the heterozygous deletion in one such BPES patient with microcephaly and short stature. They found that this patient was haploinsufficient for both *FOXL2* and *ATR* at the genomic level. They suggested that the haploinsufficiency of *ATR* may be responsible for the microcephaly and short stature observed in this patient based on the occurrence of these clinical features in ATR-defective Seckel syndrome [78]. In a complementary study, O’Driscoll and colleagues subsequently showed that cells from this BPES-*ATR*<sup>+/-</sup> patient exhibited similar cellular defects to ATR-S cells [79]. These cells failed to show significant  $\gamma$ H2AX formation and Chk1 phosphorylation following replication fork stalling. They also exhibited a similar ATR-dependent G2-M cell cycle checkpoint defect to ATR-S cells. Importantly, this phenotype was corrected following over-expression of *ATR* in the BPES-*ATR*<sup>+/-</sup> cells. This study reinforced that fact that haploinsufficiency of *ATR* has a functional impact in human cells but furthermore, that haploinsufficiency of *ATR* is associated with a human genomic disorder that exhibits microcephaly and short stature [79].

## **7. Haploinsufficiency of ATR-pathway components in other genomic disorders.**

Work from Lam and colleagues using tissue-specific conditional knockdown in mice of Chk1, an important effector kinase and substrate of ATR, showed that *Chk1* haploinsufficiency enhances mammary tumourigenesis [82]. This work suggested that an ATR-pathway component and not just *ATR* itself is sensitive to haploinsufficiency. Following the cellular characterisation of BPES- *ATR*<sup>+/-</sup>, working

on the presumption that the ATR-pathway as a whole is sensitive to a reduced gene copy number, O'Driscoll and colleagues examined ATR-pathway function in other human genomic disorders whose causative genomic deletions were known to result in haploinsufficiency of various ATR-pathway components [79].

### ***7.1 Isolated Lissencephaly Sequence (ILS) and Miller-Dieker lissencephaly syndrome (MDLS).***

Normal human brain development involves a rapid and sustained cellular proliferation originating from the rostral end of the foetal neural tube. Cerebral cortical development is achieved via a highly regulated sequence of neuroprogenitor cell division, migration and differentiation. Platelet-activating factor acetylhydrolase, isoform B1 (*PAHFAH1B1/Lis1*), located on human chromosome 17p13.3, encodes a protein, Lissencephaly 1 (Lis1), which plays a central role in neuronal migration from the neuroepithelial stem cell layer during embryonic brain development [83-85]. Mutations in or heterozygous deletions of *PAHFAH1B1/Lis1* alone causes Isolated Lissencephaly Sequence (ILS), a disorder characterised by reduced neuronal migration resulting in a cortical surface without significant invaginations, effectively a "smooth brain" (lissencephaly) (Figure 1). Larger deletions identified in some ILS patients confer a more severe grade of lissencephaly associated with additional craniofacial abnormalities (ILS+) (Figure 1). Even larger deletions extending telomerically from the *PAHFAH1B1/Lis1* gene are associated with Miller-Dieker Lissencephaly Syndrome (MDLS), which is characterised by the most severe grade of lissencephaly, craniofacial abnormalities, microcephaly and growth retardation (Figure 1) [14,86]. Unlike ILS patients, ILS+ patients and all MDLS patients are haploinsufficient for

*RPA1*, the largest subunit of the Replication Protein A complex (Figure 1) [86]. As described previously, this complex coats ssDNA generated for example at a stalled replication fork allowing the recruitment and ultimate activation of ATR [19]. Similar to what was observed using BPES-ATR<sup>+/-</sup> cells, cell lines from ILS+ and MDLS patients collectively exhibited compromised ATR-pathway function [79]. Furthermore, these cellular defects could be complemented following over-expression of *RPA1* in these cells. ILS patient cell lines with heterozygous deletions in *PAHFAH1B1/Lis1* only, by contrast, exhibited a functional ATR-pathway response (Figure 1). Therefore, ILS+ and MDLS represent two further human genomic disorders with a clinical overlap with ATR-S that also exhibit compromised ATR-pathway function at the cellular level [79].

## **7.2 Williams-Beuren syndrome (WBS)**

WBS is caused by a hemizygous sub-microscopic deletion of 1.55-1.84Mbp on chromosome 7q11.23 encompassing around 25-30 genes. The clinical presentation of this condition is multifaceted including craniofacial, endocrinological and cardiovascular abnormalities along with microcephaly and growth retardation [87,88]. One of the genes heterozygously deleted in WBS is *ELN*, which encodes elastin, a key structural component of vascular tissues. A characteristic cardiovascular abnormality of WBS, supravalvular aortic stenosis (narrowing of the ascending aorta), is thought to specifically derive from haploinsufficiency of *ELN* [89]. WBS patients are also haploinsufficient for *RFC2* a gene that encodes a component of the Replication Factor C (RFC) complex that plays a fundamental role during DNA replication loading PCNA onto chromatin facilitating DNA polymerase action [90]. *RFC2* is also a



component of the Rad17-RFC2-5 complex that is known to function in the DDR [91-94]. In fact, Rad17 is phosphorylated by ATM and ATR following DNA damage. Interestingly, a yeast strain (*S. cerevisiae*) with a hypomorphic mutation in *Rfc2* fails to activate cell cycle arrest following DNA damage suggesting that the RFC complex and specifically *RFC2* plays a direct role in the DDR [95]. O'Driscoll and colleagues also showed that WBS patient-derived cell lines exhibit a defective ATR-dependent DDR that could be complemented following re-introduction of *RFC2* into these cells [79]. Therefore, WBS represents another genomic disorder that exhibits defective ATR-pathway activity wherein microcephaly and growth retardation are included in its clinical spectrum [79].

Of course for all of the genomic disorders discussed above that exhibit a defective ATR-dependent DDR associated with microcephaly and short stature, the occurrence of this specific DDR defect with these particular clinical features is associative. This association, albeit strong, remains an association only, hence more work using complementary systems such as gene targeting or tissue-specific knockdown in the murine system will be required to definitively prove the link between ATR-pathway dysfunction and these developmental abnormalities.

## **8. Haploinsufficiency of other DDR components.**

Increasing evidence from murine gene targeting studies suggests that haploinsufficiency of various components of a diverse distinct range DDR pathways and not just *ATR* or ATR-pathway components have a functional impact on maintaining genomic stability. For example, haploinsufficiency of various

components of the mitotic spindle checkpoint such *BubR1* or *MAD2* are associated with aberrant chromosome segregation and aneuploidy in mice [96,97]. Recently, haploinsufficiency of another mitotic spindle checkpoint component, *Mad1*, has been shown to be associated with aneuploidy and increased constitutive tumour incidence in *Mad1*<sup>+/-</sup> mice compared to their wild-type (*Mad1*<sup>+/+</sup>) littermates [98]. Homologous recombination (HR) is an error-free DSB repair pathway used by mammalian cells when a homologous sister chromatid is available as a template for DNA repair (S- and G2-phases of the cell cycle) [99]. Haploinsufficiency of multiple genes whose products function in HR results in compromised genomic stability. Specific examples include *Brca1*<sup>+/-</sup>, *Xrcc2*<sup>+/-</sup>, *Xrcc3*<sup>+/-</sup>, *Rad51b*<sup>+/-</sup> and *Rad51d*<sup>+/-</sup> mice, all of which present with recombination deficiency, increased chromosomal aberrations and centrosomal fragmentation [100-105]. The Mus81-Eme1 complex functions as a structure specific endonuclease that plays a role in resolving stalling replication forks, 3'-orientated DNA flaps/overhangs and nicked HR intermediates [106]. Haploinsufficiency of both components of the structure specific endonuclease Mus81-Eme1 results in increased chromosomal aberrations and a re-replication phenotype in human and murine cells [107]. Poly(ADP-ribose) polymerase-1 (PARP-1) catalyzes the covalent attachment of long branched poly(ADP-ribose) polymers onto a diverse set of target proteins (including itself), using NAD<sup>+</sup> as its substrate. Attachment of these negatively charged polymers changes the biological activity and properties of the target proteins. PARP1 plays an important role in sensing single stranded breaks in DNA [108-110]. The level of PARP activity is sensitive to PARP-1 gene dosage [111]. Interestingly, PARP1<sup>-/+</sup> mouse embryonic fibroblasts were shown to exhibit increased supernumerary centrosomes relative to their PARP<sup>+/+</sup> counterparts [111]. Furthermore, haploinsufficiency of the histone H2A sub-family member *H2AX*, a


known ATM and ATR substrate, has been shown to compromise genomic integrity, impact on the normal response to DNA damage and enhance tumour susceptibility in the absence of *p53* in mice [112].

## **9. Implications of haploinsufficiency of DDR components in human genomic disorders.**

Murine gene targeting studies have proved invaluable in identifying DDR pathways that are sensitive to haploinsufficiency (e.g. the spindle checkpoint and HR). It is likely that haploinsufficiency of these pathways potentially contribute/will contribute to the clinical features of known and/or novel genomic disorders. A major consequence of compromised DDR is increased genomic instability and cancer predisposition (reviewed in [15]). Additionally, since DNA damaging agents are the cornerstone of clinically utilised therapeutic approaches for cancer, individuals with compromised DDR are hypersensitive to such treatments. This has been observed in Ataxia telangiectasia, Nijmegen breakage syndrome, LIG4 syndrome and Fanconi anaemia patients, some of which have fatally over-responded to standard radio- and/or chemotherapy regimens in the past (reviewed in [113]). Since increased life expectancy due to improved medical supervision is now a feature of many genomic disorders, a potentially defective DDR may become more important from the perspective of tumour development and treatment. This could be particularly relevant for genomic disorders with compromised ATR-pathway function. Whilst it is not clear whether conditions such as BPES-ATR<sup>+/-</sup>, MDLS and WBS represent tumour-predisposition conditions, it is clear that they do exhibit a defective ATR-dependent DDR [79]. Provocatively, isolated reports of malignancy in these disorders,

particularly in WBS exit [114-117]. Whether a compromised ATR-dependent DDR has a role here is currently unclear although worthy of deeper investigation.

In conclusion, plasticity of the human genome is reflected in the high level of CNV observed in clinically normal individuals. Nevertheless, CNV is associated with many pathological conditions in humans. Gene-targeting studies in mice have shown that one form of CNV, namely haploinsufficiency, of certain DDR-pathway components is associated with compromised genomic stability. Haploinsufficiency of ATR, or some of its pathway components confers a similar DDR defects to that of ATR-pathway defective Seckel syndrome cell lines. Furthermore, haploinsufficiency of *ATR*, *RPA1* and *RFC2* are associated with several human genomic disorders that exhibit microcephaly and growth retardation. Haploinsufficiency of DDR pathway components are likely to contribute to the clinical features of many genomic disorders. This will have implications for the clinical management and treatment of these conditions.



**Figure 1.** A). Chromosome 17 karyotype from a normal and MDLS individual. B). A view of a single chromatid of chromosome 17p13.3 showing the order of the genes in this region from the centromere (Cen) to the telomere (Tel). The dotted lines denote the hemizygous deletions of various sizes associated with respective patients listed on the left hand side. Con-MR is a patient with a telomeric deletion. This patient exhibits mental retardation but normal stature and head circumference. IIS represents an Isolated Lissencephaly Sequence patient deleted for *PAFAH1B1/Lis1*. IIS+ denotes Isolated Lissencephaly Sequence patients with increasing deletions telomeric from *PAFAH1B1/Lis1*. MDLS represents Miller-Dieker Lissencephaly.

Disorder	Mutant gene	ATR-dependent cellular features
Seckel syndrome	<i>ATR</i> , <i>PCNT2</i> and unknown	Defective ATR-dependent G2-M arrest, supernumerary mitotic centrosomes, DFS expression
Nijmegen breakage syndrome	<i>NBS1</i>	Defective ATR-dependent G2-M arrest
Fanconi anaemia	<i>FANC-A,B,C,D1,D2,E,F,G,H,M,J</i>	Defective ATR-dependent G2-M arrest, DFS expression
MCPH1-dependent Primary Microcephaly	<i>MCPH1</i>	Defective ATR-dependent G2-M arrest, supernumerary mitotic centrosomes

**Table 1.** Mendelian disorders that exhibit microcephaly and growth retardation associated with defective ATR-pathway function.

Disorder	Chromosome Deletion	ATR-pathway component	ATR-dependent cellular features
BPES-ATR <sup>+/-</sup>	3q23	<i>ATR</i>	Defective ATR-dependent $\gamma$ H2AX formation, Chk1 phosphorylation and G2-M arrest
Isolated Lissencephaly Sequence	17p13.3	<i>RPA1</i>	Defective ATR-dependent $\gamma$ H2AX formation, Chk1 phosphorylation and G2-M arrest
Miller-Dieker Lissencephaly Syndrome	17p13.3	<i>RPA1</i>	Defective ATR-dependent $\gamma$ H2AX formation, Chk1 phosphorylation and G2-M arrest
Williams-Beuren Syndrome	7q11.23	<i>RFC2</i>	Defective ATR-dependent G2-M arrest

**Table 2.** Genomic disorders that exhibit microcephaly and growth retardation associated with hemizygous deletions of ATR pathway components.

## Web Resources

Online Mendelian Inheritance in Man, (<http://www.ncbi.nlm.nih.gov/Omim>), for reviews of the specific genes and all of the disorders described here.

DECIPHER, (<https://decipher.sanger.ac.uk/>), a database collating multiple genomic imbalances and their associated clinical features in human genomic disorders.

EUCARUCA, (<http://agserver01.azn.nl:8080/ecaruca/whatisEc.jsp>), a database of cytogenetic and clinical data of rare chromosomal aberrations from all centres that are member of the European Cytogeneticists Association (ECA).

Database of Genomic Variants, (<http://projects.tcag.ca/variation/>), a database listing a comprehensive summary of structural variation in the human genome.

## Acknowledgements.

Special thanks Prof P.A Jeggo for all her support over the years. Thanks also to Prof. W.B Dobyns and Dr. J. M van Hagen for genomic disorder-related cell lines. M.O'D is a Cancer Research UK Senior Research Fellow whose laboratory is supported by the CRUK and UK Medical Research Council.

## References.

- [1] R.J. Roper and R.H. Reeves Understanding the Basis for Down Syndrome Phenotypes, *PLoS Genetics* 2 (2006) e50.
- [2] R. Redon, Ishikawa, Shumpei, Fitch, Karen R, Feuk, Lars Perry, George H. Andrews, T. Daniel Fiegler, Heike Shapero, Michael H. Carson, Andrew R. Chen, Wenwei Cho, Eun Kyung Dallaire, Stephanie Freeman, Jennifer L. Gonzalez, Juan R. Gratacos, Monica Huang, Jing Kalaitzopoulos, Dimitrios Komura, Daisuke MacDonald, Jeffrey R. Marshall, Christian R. Mei, Rui Montgomery, Lyndal Nishimura, Kunihiro Okamura, Kohji Shen, Fan Somerville, Martin J. Tchinda, Joelle Valsesia, Armand Woodwark, Cara Yang, Fengtang Zhang, Junjun Zerjal, Tatiana Zhang, Jane Armengol, Lluís Conrad, Donald F. Estivill, Xavier Tyler-Smith, Chris Carter, Nigel P. Aburatani, Hiroyuki Lee, Charles Jones, Keith W. Scherer, Stephen W. Hurles, Matthew E. Global variation in copy number in the human genome, *Nature* 444 (2006) 444-454.
- [3] S. Levy, G. Sutton, P.C. Ng, L. Feuk, A.L. Halpern, B.P. Walenz, N. Axelrod, J. Huang, E.F. Kirkness, G. Denisov, Y. Lin, J.R. MacDonald, A.W.C. Pang, M. Shago, T.B. Stockwell, A. Tsiamouri, V. Bafna, V. Bansal, S.A. Kravitz, D.A. Busam, K.Y. Beeson, T.C. McIntosh, K.A. Remington, J.F. Abril, J. Gill, J. Borman, Y.-H. Rogers, M.E. Frazier, S.W. Scherer, R.L. Strausberg and J.C. Venter The Diploid Genome Sequence of an Individual Human, *PLoS Biology* 5 (2007) e254.
- [4] E. Pennisi BREAKTHROUGH OF THE YEAR: Human Genetic Variation, *Science* 318 (2007) 1842-1843.
- [5] J.R. Lupski Genomic rearrangements and sporadic disease, *Nat Genet* (2007).
- [6] C. Lee, A.J. Iafrate and A.R. Brothman Copy number variations and clinical cytogenetic diagnosis of constitutional disorders, *Nat Genet* (2007).
- [7] N.P. Carter Methods and strategies for analyzing copy number variation using DNA microarrays, *Nat Genet* (2007).
- [8] J. Sebat Major changes in our DNA lead to major changes in our thinking, *Nat Genet* (2007).
- [9] S.A. McCarroll and D.M. Altshuler Copy-number variation and association studies of human disease, *Nat Genet* (2007).
- [10] G.M. Cooper, D.A. Nickerson and E.E. Eichler Mutational and selective effects on copy-number variants in the human genome, *Nat Genet* (2007).
- [11] D.F. Conrad and M.E. Hurles The population genetics of structural variation, *Nat Genet* (2007).
- [12] S.W. Scherer, C. Lee, E. Birney, D.M. Altshuler, E.E. Eichler, N.P. Carter, M.E. Hurles and L. Feuk Challenges and standards in integrating surveys of structural variation, *Nat Genet* (2007).
- [13] J.R. Lupski Genomic disorders: structural features of the genome can lead to DNA rearrangements and human disease traits, *Trends in Genetics* 14 (1998) 417-422.
- [14] J. Yingling, Toyo-Oka, K, Wynshaw-Boris, A. Miller-Dieker syndrome: analysis of a human contiguous gene syndrome in the mouse., *Am J Hum Genet* 73 (2003) 475-488.



- [15] M. O'Driscoll and P.A. Jeggo The role of double-strand break repair [mdash] insights from human genetics, *Nat Rev Genet* 7 (2006) 45-54.
- [16] R.T. Abraham Cell cycle checkpoint signaling through the ATM and ATR kinases, *Genes Dev* 15 (2001) 2177-2196.
- [17] R.T. Abraham PI 3-kinase related kinases: 'big' players in stress-induced signaling pathways, *DNA Repair* 3 (2004) 883-887.
- [18] Y. Shiloh ATM and ATR: networking cellular responses to DNA damage, *Curr Opin Genet Dev* 11 (2001) 71-77.
- [19] L. Zou, D. Liu and S.J. Elledge Replication protein A-mediated recruitment and activation of Rad17 complexes, *Proc Natl Acad Sci U S A* 100 (2003) 13827-13832.
- [20] L. Zou and S.J. Elledge Sensing DNA damage through ATRIP recognition of RPA-ssDNA complexes, *Science* 300 (2003) 1542-1548.
- [21] A. Kumagai, J. Lee, H.Y. Yoo and W.G. Dunphy TopBP1 Activates the ATR-ATRIP Complex, *Cell* 124 (2006) 943-955.
- [22] T.T. Paull, E.P. Rogakou, V. Yamazaki, C.U. Kirchgessner, M. Gellert and W.M. Bonner A critical role for histone H2AX in recruitment of repair factors to nuclear foci after DNA damage., *Curr Biol* 10 (2000) 886-895.
- [23] A. Celeste, O. Fernandez-Capetillo, M.J. Kruhlak, D.R. Pilch, D.W. Staudt, A. Lee, R.F. Bonner, W.M. Bonner and A. Nussenzweig Histone H2AX phosphorylation is dispensable for the initial recognition of DNA breaks, *Nat Cell Biol* 5 (2003) 675-679.
- [24] O. Fernandez-Capetillo, A. Lee, M. Nussenzweig and A. Nussenzweig H2AX: the histone guardian of the genome, *DNA Repair* 3 (2004) 959-967.
- [25] L. Zou, D. Cortez and S.J. Elledge Regulation of ATR substrate selection by Rad17-dependent loading of Rad9 complexes onto chromatin, *Genes Dev.* 16 (2002) 198-208.
- [26] T. Stiff, Reis, C, Alderton, G.K, Woodbine, L, O'Driscoll, M, Jeggo, P.A Nbs1 is required for ATR-dependent phosphorylation events., *EMBO J* 24 (2005) 199-208.
- [27] A. Jazayeri, J. Falck, C. Lukas, J. Bartek, G.C.M. Smith, J. Lukas and S.P. Jackson ATM- and cell cycle-dependent regulation of ATR in response to DNA double-strand breaks, *Nat Cell Biol* 8 (2006) 37-45.
- [28] T. Stiff, S. Walker, K. Cerosaletti, A. Goodarzi, E. Petermann, P. Concannon, M. O'Driscoll and P. Jeggo ATR-dependent phosphorylation and activation of ATM in response to UV treatment or replication fork stalling, *EMBO J.* 25 (2006) 5775-5782.
- [29] K. Savitsky, A. Bar-Shira, S. Gilad, G. Rotman, Y. Ziv, L. Vanagaite, D.A. Tagle, S. Smith, T. Uziel, S. Sfez, M. Ashkenazi, I. Pecker, M. Frydman, R. Harnik, S.R. Patanjali, A. Simmons, G.A. Clines, A. Sartiel, R.A. Gatti, L. Chessa, O. Sanal, M.F. Lavin, N.G.J. Jaspers, M.R. Taylor, C.F. Arlett, T. Miki, S.M. Weissman, M. Lovett, F.S. Collins and Y. Shiloh A single ataxia telangiectasia gene with a product similar to PI 3-kinase, *Science* 268 (1995) 1749-1753.
- [30] G.R.A. Chun H.H Ataxia-telangiectasia, an evolving phenotype., *DNA Repair* 3 (2004) 1187-1196.
- [31] M. O'Driscoll, V.L. Ruiz-Perez, C.G. Woods, P.A. Jeggo and J.A. Goodship A splicing mutation affecting expression of ataxia-telangiectasia and Rad3-related protein (ATR) results in Seckel syndrome, *Nature Genetics* 33 (2003) 497-501.

- [32] M. O'Driscoll and P.A. Jeggo Clinical Impact of ATR Checkpoint Signalling Failure in Humans, *Cell Cycle* 2 (2003) 194-195.
- [33] S.G. Durkin and T.W. Glover Chromosome Fragile Sites, *Annual Review of Genetics* 41 (2007) 169-192.
- [34] M.F. Arlt, S.G. Durkin, R.L. Ragland and T.W. Glover Common fragile sites as targets for chromosome rearrangements, *DNA Repair* 5 (2006) 1126-1135.
- [35] K. Mimori, T. Druck, H. Inoue, H. Alder, L. Berk, M. Mori, K. Huebner and C.M. Croce Cancer-specific chromosome alterations in the constitutive fragile region FRA3B, *Proceedings of the National Academy of Sciences* 96 (1999) 7456-7461.
- [36] H. Ishii, K. Mimori, H. Inoue, T. Inageta, K. Ishikawa, S. Semba, T. Druck, F. Trapasso, K. Tani, A. Vecchione, C.M. Croce, M. Mori and K. Huebner Fhit Modulates the DNA Damage Checkpoint Response, *Cancer Res* 66 (2006) 11287-11292.
- [37] A.M. Casper, P. Nghiem, M.F. Arlt and T.W. Glover ATR regulates fragile site stability, *Cell* 111 (2002) 779-789.
- [38] M.F. Arlt, B. Xu, S.G. Durkin, A.M. Casper, M.B. Kastan and T.W. Glover BRCA1 Is Required for Common-Fragile-Site Stability via Its G2/M Checkpoint Function, *Mol. Cell. Biol.* 24 (2004) 6701-6709.
- [39] N.G. Howlett, T. Taniguchi, S.G. Durkin, A.D. D'Andrea and T.W. Glover The Fanconi anemia pathway is required for the DNA replication stress response and for the regulation of common fragile site stability, *Hum. Mol. Genet.* 14 (2005) 693-701.
- [40] S.G. Durkin, M.F. Arlt, N.G. Howlett and T.W. Glover Depletion of CHK1, but not CHK2, induces chromosomal instability and breaks at common fragile sites, *Oncogene* 25 (2006) 4381-4388.
- [41] L.M. Pirzio, P. Pichierri, M. Bignami and A. Franchitto Werner syndrome helicase activity is essential in maintaining fragile site stability, *J. Cell Biol.* 180 (2008) 305-314.
- [42] H.P.G. Seckel Bird-headed dwarfs: Studies in developmental anthropology including human proportions, in: C.T. Thomas (Ed.), *Bird-headed dwarfs: Studies in developmental anthropology including human proportions.*, S.Karger, Basel, 1960.
- [43] J. Goodship, H. Gill, J. Carter, A. Jackson, M. Splitt and M. Wright Autozygosity mapping of a seckel syndrome locus to chromosome 3q22. 1-q24, *Am J Hum Genet* 67 (2000) 498-503.
- [44] E.J. Brown and D. Baltimore ATR disruption leads to chromosomal fragmentation and early embryonic lethality., *Genes Dev.* 14 (2000) 397-402.
- [45] A. de Klein, M. Muijtjens, R. van Os, Y. Verhoeven, B. Smit, A.M. Carr, A.R. Lehmann and J.H.J. Hoeijmakers Targeted disruption of the cell cycle checkpoint gene ATR leads to early embryonic lethality in mice, *Curr. Biol.* 10 (2000) 479-482.
- [46] G.K. Alderton, Joenje, H, Varon, R, Borglum, A D, Jeggo, P A, O'Driscoll, M. Seckel syndrome exhibits cellular features demonstrating defects in the ATR signalling pathway, *Human Molecular Genetics* 13 (2004) 3127-3138.
- [47] A.M. Casper, S.G. Durkin, M.F. Arlt and T.W. Glover Chromosomal instability at common fragile sites in seckel syndrome, *American Journal of Human Genetics* 75 (2004) 654-660.

- [48] A.D. Borglum, T. Balslev, A. Haagerup, N. Birkebaek, H. Binderup, T.A. Kruse and J.M. Hertz A new locus for Seckel syndrome on chromosome 18p11.31-q11.2, *Eur J Hum Genet* 9 (2001) 753-757.
- [49] L. Faivre, M. Le Merrer, S. Lyonnet, H. Plauchu, N. Dagoneau, A.B. Campos-Xavier, J. Attia-Sobol, A. Verloes, A. Munnich and V. Cormier-Daire Clinical and genetic heterogeneity of Seckel syndrome, *Am J Med Genet* 112 (2002) 379-383.
- [50] M.O. Kilinc, V.N. Ninis, S.A. Ugur, B. Tuysuz, M. Seven, S. Balci, J. Goodship and A. Tolun Is the novel SCKL3 at 14q23 the predominant Seckel locus?, *Eur J Hum Genet* 11 (2003) 851-857.
- [51] S.G. Hinchcliffe E.H "It takes two to tango": understanding how centrosome duplication is regulated throughout the cell cycle, *Genes Dev.* 15 (2001) 1167-1181.
- [52] K. Fukasawa Oncogenes and tumour suppressors take on centrosomes, *Nat Rev Cancer* 7 (2007) 911-924.
- [53] J.A. Golden. Cell migration and cerebral cortical development., *Neuropathol Appl Neurobiol* 27 (2001) 22-28.
- [54] J.L. Lu B, Jan Y.N. Control of cell divisions in the nervous system: symmetry and asymmetry., *Annu Rev Neurosci* 23 (2000) 531-556.
- [55] J. Bond and C.G. Woods Cytoskeletal genes regulating brain size, *Current Opinion in Cell Biology* 18 (2006) 95-101.
- [56] E. Griffith, S. Walker, C.-A. Martin, P. Vagnarelli, T. Stiff, B. Vernay, N.A. Sanna, A. Saggarr, B. Hamel, W.C. Earnshaw, P.A. Jeggo, A.P. Jackson and M. O'Driscoll Mutations in pericentrin cause Seckel syndrome with defective ATR-dependent DNA damage signaling, *Nat Genet* 40 (2008) 232-236.
- [57] A. Rauch, C.T. Thiel, D. Schindler, U. Wick, Y.J. Crow, A.B. Ekici, A.J. van Essen, T.O. Goecke, L. Al-Gazali, K.H. Chrzanowska, C. Zweier, H.G. Brunner, K. Becker, C.J. Curry, B. Dallapiccola, K. Devriendt, A. Dorfler, E. Kinning, A. Megarbane, P. Meinecke, R.K. Semple, S. Spranger, A. Toutain, R.C. Trembath, E. Voss, L. Wilson, R. Hennekam, F. de Zegher, H.-G. Dorr and A. Reis Mutations in the Pericentrin (PCNT) Gene Cause Primordial Dwarfism, *Science* 319 (2008) 816-819.
- [58] B. Delaval and S. Doxsey GENETICS: Dwarfism, Where Pericentrin Gains Stature, *Science* 319 (2008) 732-733.
- [59] R. Varon, C. Vissinga, M. Platzer, K.M. Cerosaletti, K.H. Chrzanowska, K. Saar, G. Beckmann, E. Seemanova, P.R. Cooper, N.J. Nowak, M. Stumm, C.M.R. Weemaes, R.A. Gatti, R.K. Wilson, M. Digweed, A. Rosenthal, K. Sperling, P. Concannon and A. Reis Nibrin, a novel DNA double-strand break repair protein, is mutated in Nijmegen breakage syndrome, *Cell* 93 (1998) 467-476.
- [60] S.K. Digweed M Nijmegen breakage syndrome: clinical manifestation of defective response to DNA double-strand breaks., *DNA Repair* 3 (2004) 1207-1217.
- [61] J.H. Lee and T.T. Paull Direct activation of the ATM protein kinase by the Mre11/Rad50/Nbs1 complex, *Science* 304 (2004) 93-96.
- [62] Wang X and D.A. AD. The interplay of Fanconi anemia proteins in the DNA damage response., *DNA Repair* 3 (2004) 1063-1069.
- [63] K.J. Patel and H. Joenje Fanconi anemia and DNA replication repair, *DNA Repair* 6 (2007) 885-890.

- [64] P.R. Andreassen, A.D. D'Andrea and T. Taniguchi ATR couples FANCD2 monoubiquitination to the DNA damage response, *Genes and Development* 18 (2004) 1958-1963.
- [65] R.C. Gregory, T. Taniguchi and A.D. D'Andrea Regulation of the Fanconi anemia pathway by monoubiquitination, *Semin Cancer Biol* 13 (2003) 77-82.
- [66] K. Nakanishi, T. Taniguchi, V. Ranganathan, H. New, V. , L. Moreau, A. , M. Stotsky, C. Mathew, G. , M. Kastan, B. , D. Weave, T. and A. D'Andrea, D. Interaction of FANCD2 and NBS1 in the DNA damage response, *Nat Cell Biol* 4 (2002) 913-920.
- [67] G.-H.I. Taniguchi T, Xu B, Andreassen PR, Gregory RC, Kim ST, Lane WS, Kastan MB, D'Andrea AD Convergence of the fanconi anemia and ataxia telangiectasia signaling pathways., *Cell* 4 (2002) 459-472.
- [68] Bond J, Roberts E, Springell K, Lizarraga SB, Scott S, Higgins J, Hampshire DJ, Morrison EE, Leal GF, Silva EO, Costa SM, Baralle D, Raponi M, Karbani G, Rashid Y, Jafri H, Bennett C, Corry P, Walsh CA and W. CG. A centrosomal mechanism involving CDK5RAP2 and CENPJ controls brain size., *Nat Genet* 37 (2005) 353-355.
- [69] C.G. Woods Human microcephaly., *Curr Opin Neurobiol* 14 (2004) 112-117.
- [70] R.E. Bond J, Mochida G. H, Hampshire D. J, Scott S, Askham J. M, Springell K, Mahadevan M, Crow Y. J, Markham A. F, Walsh C. A, Woods C. G. ASPM is a major determinant of cerebral cortical size, *Nat Genet* 32 (2002) 316-320.
- [71] A.P. Jackson, H. Eastwood, S.M. Bell, J. Adu, C. Toomes, I.M. Carr, E. Roberts, D.J. Hampshire, Y.J. Crow, A.J. Mighell, G. Karbani, H. Jafri, Y. Rashid, R.F. Mueller, A.F. Markham and C.G. Woods Identification of microcephalin, a protein implicated in determining the size of the human brain, *Am J Hum Genet* 71 (2002) 136-142.
- [72] S.-Y. Lin, R. Rai, K. Li, Z.-X. Xu and S.J. Elledge BRIT1/MCPH1 is a DNA damage responsive protein that regulates the Brca1-Chk1 pathway, implicating checkpoint dysfunction in microcephaly, *PNAS* 102 (2005) 15105-15109.
- [73] X. Xu, J. Lee and D.F. Stern Microcephalin Is a DNA Damage Response Protein Involved in Regulation of CHK1 and BRCA1, *J. Biol. Chem.* 279 (2004) 34091-34094.
- [74] G.K. Alderton, Galbiati, L, Griffith, E, Surinya, K.H, Neitzel, H, Jackson, A.P, Jeggo, P.A, O'Driscoll, M. Regulation of mitotic entry by microcephalin and its overlap with ATR signalling, *Nature Cell Biology* 8 (2006) 725-733.
- [75] A.L. Brown, C.H. Lee, J.K. Schwarz, N. Mitiku, H. Piwnica-Worms and J.H. Chung A human Cds1-related kinase that functions downstream of ATM protein in the cellular response to DNA damage, *Proc Natl Acad Sci U S A* 96 (1999) 3745-3750.
- [76] K.A. Lewis, Mullany, Sally, Thomas, Bijoy, Chien, Jeremy, Loewen, Ralitsa, Shridhar, Viji, Cliby, William A. Heterozygous ATR Mutations in Mismatch Repair-Deficient Cancer Cells Have Functional Significance, *Cancer Res* 65 (2005) 7091-7095.
- [77] Y. Fang, Tsao, Cheng-Chung, Goodman, Barbara K, Furumai, Ryohei, Tirado, Carlos A, Abraham, Robert T, Wang, Xiao-Fan. ATR functions as a gene dosage-dependent tumor suppressor on a mismatch repair-deficient background, *The EMBO Journal* 23 (2004) 3164-3174.

- [78] M. de Ru, Gille, JJ, Nieuwint, AWM, Bijlsma, JJ, van der Blij, JF, van Hagen, JM. Interstitial deletion in 3q in a patient with blepharophimosis-ptosis-epicanthus inversus syndrome (BPES) and microcephaly, mild mental retardation and growth delay: Clinical report and review of the literature, *American Journal of Medical Genetics Part A* 137A (2005) 81-87.
- [79] M. O' Driscoll, W.B. Dobyns, J.M. van Hagen and P.A. Jeggo Cellular and Clinical Impact of Haploinsufficiency for Genes Involved in ATR Signaling, *American journal of human genetics* 81 (2007) 77-86.
- [80] L. Crisponi, Deiana, Manila, Loi, Angela, Chiappe, Francesca Uda, Manuela Amati, Patrizia Bisceglia, Luigi Zelante, Leopoldo Nagaraja, Ramaiah Porcu, Susanna Serafina Ristaldi, M. Marzella, Rosalia Rocchi, Mariano Nicolino, Marc, Lienhardt-Roussie, Nivelon, Anne, Verloes, Annie, Schlessinger, Alain, Gasparini, David, Bonneau, Paolo, Cao, Dominique, Pilia, Antonio, Giuseppe The putative forkhead transcription factor FOXL2 is mutated in blepharophimosis/ptosis/epicanthus inversus syndrome, *Nat Genet* 27 (2001) 159-166.
- [81] E. De Baere, Dixon, Michael J, Small, Kent W. Jabs, Ethylin W. Leroy, Bart P. Devriendt, Koenraad Gillerot, Yves Mortier, Geert Meire, Francoise Van Maldergem, Lionel Courtens, Winnie Hjalgrim, Helle Huang, Shangzhi, I.V.R. Liebaers, Nicole Touraine, Philippe Praphanphoj, Verayuth Verloes, Alain Udar, Nitin and V.C. Yellore, Meenal Yelchits, Svetlana De Paepe, Anne Kuttann, Frederique Fellous, Marc Veitia, Reiner Messiaen, Ludwine Spectrum of FOXL2 gene mutations in blepharophimosis-ptosis-epicanthus inversus (BPES) families demonstrates a genotype-phenotype correlation, *Hum. Mol. Genet.* 10 (2001) 1591-1600.
- [82] M.H. Lam, Q. Liu, S.J. Elledge and J.M. Rosen Chk1 is haploinsufficient for multiple functions critical to tumor suppression, *Cancer Cell* 6 (2004) 45-59.
- [83] C. Cardoso, Leventer R.J, Matsumoto N, Kuc J.A, Ramocki M.B, Mewborn S.K, Dudliceck L.L, May L.F, Mills P.L, Das S, Pilz D.T, Dobyns W.B, Ledbetter D.H. The location and type of mutation predict malformation severity in isolated lissencephaly caused by abnormalities within the LIS1 gene., *Hum Mol Genet* 9 (2000) 3019-3028.
- [84] R.J. Leventer, Cardoso, C, Ledbetter, D.H, Dobyns, W.B. LIS1: from cortical malformation to essential protein of cellular dynamics., *Trends Neurosci* 24 (2001) 489-492.
- [85] C. Cardoso, Leventer, R.J, Dowling, J.J, Ward, H.L, Chung, J, Petras, K.S, Roseberry, J.A, Weiss, A.M, Das, S, Martin, C.L, Pilz, D.T, Dobyns, W.B, Ledbetter, D.H. Clinical and molecular basis of classical lissencephaly: Mutations in the LIS1 gene (PAFAH1B1). *Hum Mutat* 19 (2002) 4-15.
- [86] C. Cardoso, Leventer, R.J, Ward, H.L, Toyo-Oka, K, Chung, J, Gross, A, Martin, C.L, Allanson, J, Pilz, D.T, and A. Olney, H. Mutchinick, O.M. Hirotsune, S. Wynshaw-Boris, A. Dobyns, W.B. Ledbetter, D.H. Refinement of a 400-kb critical region allows genotypic differentiation between isolated lissencephaly, Miller-Dieker syndrome, and other phenotypes secondary to deletions of 17p13.3., *Am J Hum Genet* 72 (2003) 918-930.
- [87] M. Tassabehji Williams-Beuren syndrome: a challenge for genotype-phenotype correlations., *Hum Mol Genet.* 12 (2003) 229-237.
- [88] Y.Q. Wu, Sutton, V.R, Nickerson, E, Lupski, J.R, Potocki, L, Korenberg, J.R, Greenberg, F, Tassabehji, M, Shaffer, L.G. Delineation of the common critical

- region in Williams syndrome and clinical correlation of growth, heart defects, ethnicity, and parental origin., *Am J Med Genet.* 78 (1998) 82-89.
- [89] M. Tassabehji, K. Metcalfe, A. Karmiloff-Smith, M. Carette, J. , J. Grant, N. Dennis, W. Reardon, M. Splitt, A. Read, P. and D. Donnai Williams syndrome: use of chromosomal microdeletions as a tool to dissect cognitive and physical phenotypes., *Am J Hum Genet* 64 (1999) 118-125.
  - [90] V. Ellison and B. Stillman Reconstitution of recombinant human replication factor C (RFC) and identification of an RFC subcomplex possessing DNA-dependent ATPase activity, *J. Biol. Chem.* 273 (1998) 5979-5987.
  - [91] J. Griffith, L.A. Lindsey-Boltz and A. Sancar Structures of the human Rad17-replication factor C and checkpoint Rad 9-1-1 complexes visualized by glycerol spray/low voltage microscopy., *J .Biol. Chem.* 277 (2002) 15233-15236.
  - [92] V.P. Bermudez, L.A. Lindsey-Boltz, A.J. Cesare, Y. Maniwa, J.D. Griffith, J. Hurwitz and A. Sancar Loading of the human 9-1-1 checkpoint complex onto DNA by the checkpoint clamp loader hRad17-replication factor C complex in vitro., *Proc Natl Acad Sci U S A.* 100 (2003) 1633-1638.
  - [93] Z.L. Wang X, Zheng H, Wei Q, Elledge S.J, Li L. Genomic instability and endoreduplication triggered by RAD17 deletion., *Genes Dev.* 17 (2003) 965-970.
  - [94] L. Zou, D. Liu and S.J. Elledge Replication protein A-mediated recruitment and activation of Rad17 complexes, *PNAS* 100 (2003) 13827-13832.
  - [95] V.N. Noskov, Araki, H, Sugino, A. The RFC2 gene, encoding the third-largest subunit of the replication factor C complex, is required for an S-phase checkpoint in *Saccharomyces cerevisiae*, *Mol Cell Biol.* 18 (1998) 4914-4923.
  - [96] D. Baker, J. , K. Jeganathan, B. , J. Cameron, D. , M. Thompson, S. Juneja, A. Kopecka, R. Kumar, R. Jenkins, B. , P. de Groen, C. , P. Roche and J.M. van Deursen BubR1 insufficiency causes early onset of aging-associated phenotypes and infertility in mice., *Nat Genet.* 2004 36 (2004) 744-749.
  - [97] L. Michel, S. , V. Liberal, A. Chatterjee, R. Kirchwegger, B. Pasche, W. Gerald, M. Dobles, P. Sorger, K. , V. Murty, V. and R. Benezra MAD2 haplo-insufficiency causes premature anaphase and chromosome instability in mammalian cells, *Nature.* 409 (2001) 355-359.
  - [98] Y. Iwanaga, Y.-H. Chi, A. Miyazato, S. Sheleg, K. Haller, J.-M. Peloponese, Jr., Y. Li, J.M. Ward, R. Benezra and K.-T. Jeang Heterozygous Deletion of Mitotic Arrest-Deficient Protein 1 (MAD1) Increases the Incidence of Tumors in Mice, *Cancer Res* 67 (2007) 160-166.
  - [99] T. Helleday Pathways for mitotic homologous recombination in mammalian cells, *Mutat Res* 532 (2003) 103-115.
  - [100] X. Xu, Z. Weaver, S.P. Linke, C. Li, J. Gotay, X.W. Wang, C.C. Harris, T. Ried and C.X. Deng Centrosome amplification and a defective G2-M cell cycle checkpoint induce genetic instability in BRCA1 exon 11 isoform-deficient cells., *Mol Cell* 3 (1999) 389-395.
  - [101] B. Deans, C.S. Griffin, M. Maconochie and J. Thacker Xrcc2 is required for genetic stability, embryonic neurogenesis and viability in mice, *Embo J* 19 (2000) 6675-6685.
  - [102] C. Griffin, S. , P. Simpson, J. , C. Wilson, R. and J. Thacker Mammalian recombination-repair genes XRCC2 and XRCC3 promote correct chromosome segregation, *Nat Cell Biol* 2 (2000) 757-761.

- [103] B. Deans, C.S. Griffin, P. O'Regan, M. Jasin and J. Thacker Homologous recombination deficiency leads to profound genetic instability in cells derived from Xrcc2-knockout mice, *Cancer Res* 63 (2003) 8181-8187.
- [104] O. Date, M. Katsura, M. Ishida, T. Yoshihara, A. Kinomura, T. Sueda and K. Miyagawa Haploinsufficiency of RAD51B Causes Centrosome Fragmentation and Aneuploidy in Human Cells, *Cancer Res* 66 (2006) 6018-6024.
- [105] P.G. Smiraldo, A.M. Gruver, J.C. Osborn and D.L. Pittman Extensive Chromosomal Instability in Rad51d-Deficient Mouse Cells, *Cancer Res* 65 (2005) 2089-2096.
- [106] F. Osman and M.C. Whitby Exploring the roles of Mus81-Eme1/Mms4 at perturbed replication forks, *DNA Repair* 6 (2007) 1004-1017.
- [107] T. Hiyama, M. Katsura, T. Yoshihara, M. Ishida, A. Kinomura, T. Tonda, T. Asahara and K. Miyagawa Haploinsufficiency of the Mus81-Eme1 endonuclease activates the intra-S-phase and G2/M checkpoints and promotes rereplication in human cells, *Nucl. Acids Res.* 34 (2006) 880-892.
- [108] P.A. Jeggo DNA repair: PARP - another guardian angel?, *Current Biology* 8 (1998) 49-51.
- [109] S.F. El-Khamisy, M. Masutani, H. Suzuki and K.W. Caldecott A requirement for PARP-1 for the assembly or stability of XRCC1 nuclear foci at sites of oxidative DNA damage, *Nucleic Acids Res* 31 (2003) 5526-5533.
- [110] V. Schreiber, F.o. Dantzer, J.-C. Ame and G. de Murcia Poly(ADP-ribose): novel functions for an old molecule, *Nat Rev Mol Cell Biol* 7 (2006) 517-528.
- [111] M. Kanai, W.-M. Tong, Z.-Q. Wang and M. Miwa Haploinsufficiency of poly(ADP-ribose) polymerase-1-mediated poly(ADP-ribosyl)ation for centrosome duplication, *Biochemical and Biophysical Research Communications* 359 (2007) 426-430.
- [112] A. Celeste, S. Difilippantonio, M. Difilippantonio, J. , O. Fernandez-Capetillo, D. Pilch, R., O. Sedelnikova, A. , M. Eckhaus, T. Ried, W. Bonner, M. and A. Nussenzweig H2AX haploinsufficiency modifies genomic stability and tumor susceptibility., *Cell*. 114 (2003) 371-383.
- [113] M. O'Driscoll and P.A. Jeggo CsA can induce DNA double-strand breaks: implications for BMT regimens particularly for individuals with defective DNA repair, *Bone Marrow Transplant* (2008).
- [114] U. Shizuyo, K. Masaru, K. Shigekazu and W. Michihiko Gallbladder cancer in a patient with Miller-Dieker syndrome, *Acta Paediatrica* 95 (2006) 113-114.
- [115] S. Amenta, Moschovi, Maria, Sofocleous, Christalena, Kostaridou, Stavroula, Mavrou, Ariadni, Fryssira, Helen. Non-Hodgkin lymphoma in a child with Williams syndrome, *Cancer Genetics and Cytogenetics* 154 (2004) 86-88.
- [116] Thornburg C, Roulston D and V. Castle Burkitt lymphoma and Williams syndrome: a model for children with a multisystem disorder and malignancy., *J Pediatr Hematol Oncol* 27 (2005 ) 109-111.
- [117] Semmekrot BA, Rotteveel JJ, Bakker-Niezen SH and L. F. Occurrence of an astrocytoma in a patient with Williams syndrome., *Pediatr Neurosci* 12 (1985-1986) 188-191.